arXiv:astro-ph/0301310v1 16 Jan 2003

# Limits on the Dark Energy Parameters from Cosmic Microwave Background experiments

Dmitri Pogosyan[1], J. Richard Bond[2], Carlo Contaldi[2]

[1] *Physics Department, University of Alberta, Edmonton, AB, T6G 2J1, Canada*
[2] *CITA, University of Toronto, Toronto, ON, M5S 3H8, Canada*

**Abstract.** Full suite of the present day Cosmic Microwave background (CMB) data, when combined with weak prior information on the Hubble constant and the age of the Universe, or the Large-Scale structure, provides strong indication for a non-zero density of the vacuum-like dark energy in our universe. This result independently supports the conclusions from Supernovae Ia (SN1a) data. When the model parameter space is extended to allow for the range of the equation of state parameter $w_Q$ for the dynamical field $Q$ which gives rise to dark energy, the CMB data is found to give a weak upper bound $w_Q < -0.4$ at 95% CL, however combined with SN1a data it strongly favours $w_Q < -0.8$, consistent with $\Lambda$-term like behaviour.

Measurements of anisotropies in the cosmic microwave background (CMB) radiation now tightly constrain the nature and composition of our universe. High signal-to-noise detections of primordial anisotropies have been made at angular scales ranging from the quadrupole [2] to as small as several arcminutes [11, 13, 5]. Within the context of models with adiabatic initial perturbations, as are generally predicted by inflation, these measurements have been used in combination with various other cosmological constraints to estimate the values of many important cosmological parameters. Combining their CMB data with weak cosmological constraints such as a very loose prior on the Hubble constant $H_0$, various teams have made robust determinations of several parameters, including the total energy density of the universe $\Omega_{tot}$, the density of baryons $\omega_b = \Omega_b h^2, h = H_0/100\,\mathrm{k}m/\mathrm{s}/\mathrm{M}pc$, and $n_s$, the spectral index of primordial perturbations [9, 1, 15, 12].

In Table 1 we present some of our results for the cosmological parameters together with 68% CL errors, obtained from the latest suite of CMB observations. Full details can be found in [16, 6]. Our analysis has included Acbar [8],Archeops [3], Boomerang [16], CBI [13], DASI [7], MAXIMA [10], VSA [17] and COBE-DMR [2] data. Combining these highly consistent datasets allows for a very tight determination of the baryon and dark matter physical densities $\omega_b$ and $\omega_{cdm}$ and the slope of scalar perturbations $n_s$, even with no prior restrictions on the parameters.

**Table 1.** Cosmological parameters from combination of CMB experiments

| Priors | $\Omega_{tot}$ | $n_s$ | $\omega_b$ | $\omega_{cdm}$ | $\Omega_\Lambda$ | $\tau_c$ |
|---|---|---|---|---|---|---|
| no prior | $1.2^{0.11}_{0.10}$ | $0.96^{0.04}_{0.04}$ | $0.022^{0.002}_{0.002}$ | $0.11^{0.02}_{0.02}$ | $0.24^{0.21}_{0.15}$ | $< 0.33$ |
| wk | $1.06^{0.04}_{0.04}$ | $0.95^{0.04}_{0.03}$ | $0.022^{0.002}_{0.002}$ | $0.12^{0.02}_{0.02}$ | $0.56^{0.13}_{0.14}$ | $< 0.31$ |
| wk+LSS | $1.05^{0.03}_{0.03}$ | $0.96^{0.03}_{0.03}$ | $0.022^{0.002}_{0.002}$ | $0.11^{0.02}_{0.02}$ | $0.66^{0.09}_{0.09}$ | $< 0.27$ |
| wk+SN | $1.03^{0.03}_{0.03}$ | $0.96^{0.03}_{0.03}$ | $0.023^{0.002}_{0.002}$ | $0.10^{0.02}_{0.01}$ | $0.70^{0.04}_{0.05}$ | $< 0.39$ |
| wk+LSS+SN | $1.03^{0.03}_{0.03}$ | $0.97^{0.03}_{0.03}$ | $0.023^{0.002}_{0.002}$ | $0.10^{0.02}_{0.01}$ | $0.70^{0.04}_{0.04}$ | $< 0.37$ |
| $\Omega_{tot} = 1$ plus | | | | | | |
| wk | (1.00) | $0.96^{0.03}_{0.03}$ | $0.022^{0.002}_{0.002}$ | $0.12^{0.02}_{0.02}$ | $0.72^{0.07}_{0.08}$ | $< 0.17$ |
| wk+LSS | (1.00) | $0.97^{0.03}_{0.03}$ | $0.022^{0.002}_{0.002}$ | $0.12^{0.01}_{0.01}$ | $0.70^{0.05}_{0.04}$ | $< 0.17$ |
| wk+SN | (1.00) | $0.96^{0.03}_{0.03}$ | $0.022^{0.002}_{0.002}$ | $0.12^{0.01}_{0.01}$ | $0.71^{0.05}_{0.04}$ | $< 0.18$ |
| wk+LSS+SN | (1.00) | $0.97^{0.02}_{0.03}$ | $0.022^{0.002}_{0.002}$ | $0.12^{0.01}_{0.01}$ | $0.70^{0.04}_{0.04}$ | $< 0.18$ |

Here we shall focus on the parameters describing the vacuum (or dark) energy, which may dominate the present day energy balance. In the minimal parameter set of Table 1 the vacuum energy is assumed to behave as $\Lambda$-term being described just by its density parameter $\Omega_\Lambda$. Supernovae Ia results [14] provide a strong indication that $\Omega_\Lambda > 0$.

The non-prior CMB analysis exhibits persistent degeneracy in the $\Omega_{tot} - \Omega_\Lambda$ space (see Figure 1), which extends along the direction of the constant angular size of the sound horizon at recombination — from closed models $\Omega_{tot} \sim 1.2$ with little if any vacuum energy contribution, to flat models $\Omega_{tot} \approx 1$, $\Omega_\Lambda \approx 0.7$. The degeneracy is reduced and the high likelihood region is concentrated near flat models with significant vacuum energy as soon as even weak prior restrictions ("wk" in Table 1) are imposed on the Hubble constant $0.45 < h < 0.9$ and the age of the universe $t > 10\,\mathrm{Gyr}$, or when CMB data is combined with the Large-Scale Structure (LSS) information, namely the amplitude $\sigma_8$ and the slope of density power spectrum on galaxy cluster scale (see [6] for the details of the LSS prior). Thus, present day CMB data provides new independent indication for non zero vacuum density. This conclusion is significantly strengthened by the new data obtained this year, in particular, by the Archeops team which has measured CMB temperature fluctuations in the intermediate angular range $l \sim 20 - 150$, only weakly constrained by previous data.

We note that the data indicates that the optical depth due to late-time reionization in the Universe is not large, with 95% upper limit being in the range $\tau_c < 0.3 - 0.4$. The exact value, however, is not yet robust.

The plausible interpretation of the dark energy is that it is the energy of a "quintessence" scalar field $Q$, dynamical in origin, thus, $\Omega_\Lambda = \Omega_Q$. A popular phenomenology is to describe additional complexity by one more parameter



Figure 1: Contour regions of the marginalized likelihood in $\Omega_m = 1 - \Omega_{tot} - \Omega_\Lambda$ versus $\Omega_\Lambda$ plane. Results from CMB data (blue) are superimposed on the contours from SN1a data (orange). Shaded areas correspond to 68% and 95% confidence regions in the Gaussian limit. Degeneracy of the likelihood along the line of the constant angular size of the sound horizon exhibited by CMB data (upper left) is reduced by either imposing a weak prior on the Hubble constant and age (bottom left) or by the Large Scale Structure restrictions (upper right) or both (bottom right).

$w_Q = P_Q/\rho_Q$, the ratio of the pressure $P_Q$ and the density $\rho_Q$ of the field which serves as an effective equation of state. As long as we do not have a pure $\Lambda$ term with $w_Q = -1$, the $w_Q$ is expected to vary in time, but the precision of CMB data does not yet warrant such detailed parametrization. We consider $w_Q$ constant and restict our attention to inflation motivated flat $\Omega_{tot} = 1$ cosmological models (this prior does not affect other parameters, cf. lower part of Table 1). As another simplification we neglect the effect of late-time perturbations in the $Q$ field, which may affect CMB signal at the largest scales. This effect is dependent in detail on the exact model for quintessence potential (one should be cautioned not to extend the equation of state analogy too far, and treat perturbations in the $Q$ component hydrodynamically).

Figure 2 shows that CMB alone does not yield useful restriction on the $w_Q$



Figure 2: Contours of marginalized likelihood for the parameters of the $Q$-field $\Omega_Q, w_Q$ in the class of spatially flat $\Omega_{tot} = 1$ cosmological models. Left – CMB with weak prior, right – joint CMB and SN1a results (blue) are superimposed on SN1a limits (orange). Shaded regions correspond to $68\%, 95\%$ (the lightest colour) and $99\%$ CL limits in the Gaussian approximation.

($w_Q < 0.4$ at 95% CL) but its combination with SN1a data prefers at 95% CL the low values $w_Q < -0.8$, an improvement to the limit reported in [4].

**Acknowledgements.** We are grateful to Boomerang, CBI and Acbar collaborations with whom much of the work reviewed here has been done. The computational facilities at Toronto are funded by the Canadian Fund for Innovation.

## References

[1] Balbi A., et al., 2000, ApJ 545, L1.
[2] Bennett C. L., et al. 1996, ApJ 464, L1.
[3] Benoit A., et al., 2002, submitted to A&A Letter, `astro-ph/0210305`.
[4] Bond J.R., et al., 2002, eds. V. Elias, R. Epp, R. Myers, in *Theoretical Physics, MRST 2002: A Tribute to George Libbrandt*, AIP Conf. Proceedings 646, 15-33.
[5] Dawson K. S., Holzapfel W. L., Carlstrom J. E., LaRoque S. J., Miller A. D., Nagai D., & Joy M., 2002, American Astronomical Society Meeting, 200.
[6] Goldstein J. H., et al., 2002, submitted to ApJ , `astro-ph/0212517`.
[7] Halverson, N. W., et al., 2001, ApJL , preprint, `astro-ph/0104489`.
[8] Kuo C. L., et al., 2002, submitted to ApJ , `astro-ph/0212289`.
[9] Lange A. E., et al. 2001, PRD 63, 042001.
[10] Lee A. T., et al., 2001, ApJ 561, L1-L6.
[11] Mason B. S., et al., 2002, submitted to ApJ , `astro-ph/0205384`.
[12] Netterfield C. B., et al., 2002, ApJ 571, 604.
[13] Pearson T. J., et al., 2002, submitted to ApJ , `astro-ph/0205388`.
[14] Perlmutter S., et al., 1999, ApJ 517, 565-586.
[15] Pryke C., Halverson N. W., Leitch E. M., Kovac J., & Carlstrom J. E., 2001, ApJL , preprint `astro-ph/0104490`
[16] Ruhl J. E., et al., 2002, submitted to ApJ , `astro-ph/0212229`.
[17] Scott P. F, et al., 2002, Submitted to MNRAS , `astro-ph/0205380`.